

Bose-Einstein Condensation Picture of Superconductivity in $\text{YBa}_2\text{Cu}_3\text{O}_7(91\text{K})$ and $\text{YBa}_2\text{Cu}_3\text{Se}_7(371\text{K})$. (Dilute metals).

V. N. Bogomolov

*A. F. Ioffe Physical & Technical Institute,
Russian Academy of Science,
194021 St. Petersburg, Russia**

(Dated: February 2, 2008)

A metal dilution degree in the compounds $\text{YBa}_2\text{Cu}_3\text{O}_7$ and $\text{YBa}_2\text{Cu}_3\text{Se}_7$ is defined as $z = (r_{\text{Me}}/r_{\text{O}^{2-};\text{Se}^{2-}})^3$. A substitution of oxygen by selenium changes z by 8 times and the Bose-Einstein condensation temperature equals $T_{\text{cSe}} = T_{\text{cO}} \cdot 8^{2/3} = 364\text{ K}$. The "active" electron pairs density in $\text{YBa}_2\text{Cu}_3\text{O}_7$ is about $1.7 \times 10^{20}\text{ cm}^{-3}$. The electron effective mass is about $5m_e$ and is proportional to the dielectric constant.

PACS numbers: 71.30.+h, 74.20.-z, 74.25.Jb

The superconductivity at 371 K in $\text{YBa}_2\text{Cu}_3\text{O}_7$ was found measuring the magnetic properties in [1]. However, the substitution of the oxygen ions ($r_{\text{O}^{2-}} \sim 0.5\text{\AA}$) by the selenium ones ($r_{\text{Se}^{2-}} \sim 1.0\text{\AA}$) made the lattice unstable and the compound irreproducible.

Densities of the electron pairs, transition temperatures and effective masses at the "physical" dilution of metals in $\text{Na}_{0.04}\text{NH}_3$ ($T_c \sim 200\text{ K}$, $m^* \sim 5m_e$) [2], in $\text{Na}_{0.05}\text{WO}_3$ ($T_c \sim 91\text{ K}$, $m^* \sim 10m_e$) [3], and in $\text{Ag}_2(\text{Ag}_3\text{Pb}_2\text{H}_2\text{O}_6)$ ($T_c \sim 400\text{ K}$, $m^* \sim 7.5m_e$) [4] are known and comply with the BEC model [5].

The "active" electron pair density in oxides of the $(1-2-3-7)$ type corresponds to the metal orbital occupancy, which depends upon the acceptor properties of the oxygen or selenium ions (the "chemical" dilution) [6]. (The "active" electron density in the Noble Gases condensates determines by interaction of the atom ground state orbitals [6]).

Let us assume that the Me and O^{2-} orbital occupancy are inversely proportional to the orbital volumes: $z = (r_{\text{Me}}/r_{\text{O}^{2-}})^3$; $z = 8.0$ for the pair $\text{Se}^{2-} - -\text{O}^{2-}$.

*Electronic address: V.Bogomolov@mail.ioffe.ru

Therefore, the pair density in YBaCuSe is 8 times larger, than in YBaCuO . For the BEC mechanism, $T_{\text{cSe}} = T_{\text{cO}} \cdot 8^{2/3} = 91 \cdot 4 = 364\text{ K}$ that complies with the data [1].

About one metal atom falls at one oxygen atom in YBaCuO . The pair number in the unit cell equals 2.5. Every electron pair occupies the volume $69.4\text{\AA}^3$. For the occupancy $z_{\text{Ba-O}} = (2.19/0.5)^3 = 84$ we have for the "active" electron pair density is about $1.7 \times 10^{20}\text{ cm}^{-3}$. The transition temperature $T_{\text{cO}} = 92\text{ K}$ for the BEC mechanism and the effective mass $m^* \sim 5m_e$ (the dielectric constant ~ 20).

The effective electron mass magnitudes are very close in all of these four systems [1, 2, 3, 4] and comply with the dielectric constants.

The compounds studied in [1] and [4] show the superconductivity of the "gossamer" kind.

A use of hydrogen for blocking of a part of the oxygen valence can preserve the lattice structure at the electron concentration increase leading to an increase of T_{cO} [7].

-
- [1] V.D. Shabetnik, S.Yu. Butuzov, V.I. Plaksii, *Tech. Phys. Lett.*, **21**, 382 (1995).
 - [2] R.A. Ogg, Jr., *Phys. Rev.* **69**, 243 (1946).
 - [3] A. Shengelaya, S. Reich, Y. Tsabba and K.A. Muller, *Eur. Phys. J. B* **12**, 13 (1999).
 - [4] Djurek, Z. Medunic, A.Tonejc, M. Paljevic, *Physica*

- C341-348**, 723-725 (2000); e-print: cond-mat/0310011.
- [5] V.N. Bogomolov, e-print: cond-mat/0505718 .
- [6] V.N. Bogomolov, e-print: cond-mat/0302034; /0311265; /0406564; /0412740.
- [7] V.N. Bogomolov, e-print: cond-mat/0411574.